

COLLIDING WINDS IN THE STELLAR CORE AT THE GALACTIC CENTER: SOME IMPLICATIONS

Leonid M. Ozernoy^{1,2}, Reinhard Genzel³, and Vladimir V. Usov⁴

¹ Computational Sciences Institute and Department of Physics & Astronomy, George Mason U., Fairfax, VA 22030-4444, USA

² Laboratory for Astronomy and Solar Physics, Goddard Space Flight Center, Greenbelt, MD 20771, USA; e-mail: ozernoy@hubble.gmu.edu, ozernoy@stars.gsfc.nasa.gov

³ Max-Planck-Institut für Extraterrestrische Physik, Postfach 1603, Garching bei München, 85740 Germany; e-mail: genzel@mpe.mpe-garching.mpg.de

⁴ Department of Physics, Weizmann Institute, Rehovot 76100, Israel; e-mail: fnusov@weizmann.weizmann.ac.il

ABSTRACT

We point out that a high number density of stars in the core of a dense star cluster such as the central stellar cluster at the Galactic center, where many stars possess strong stellar winds, should result in collisions of those winds. The wind collisions in the dense stellar core at the Galactic center would result in production of strong X-ray flares with the rate of $\sim 10^{-4} (N_w/10^3)^2 \text{ yr}^{-1}$ and duration of ~ 1 week, where N_w is the number of the wind producing stars in the core. Presence of a massive black hole would enhance the stellar density around it and would make collisions of the winds in the core substantially more frequent. Collisions of the stellar winds in the cluster have also a number of interesting observable implications, including generation of γ -rays by particles accelerated by the shocks from the colliding winds. These processes are also expected to be relevant to compact regions of intense star formation elsewhere.

Key words: Galaxy: center – galaxies: nuclei – infrared: – stars: early-type – stars: mass loss – X-rays: stars – shock waves – particle acceleration – gamma rays.

1. INTRODUCTION

The first near-IR observations of a broad HeI emission line from extended source IRS 16 at the Galactic center have indicated an intense outflow of matter from it (Hall et al. 1982; Geballe et al. 1982). Subsequent detection, in addition to the above HeI line, of broad Br α and Br γ HI emission lines, basically confirmed that interpretation (Geballe et al. 1984). Detailed near-IR observations of the Galactic center indicate that the broad line region as a source of mass outflow has the following approximate parameters:

$$\begin{aligned} \text{characteristic linear size (FWHM)} \quad l &\approx 3'' \approx 0.1 \text{ pc}, \\ \text{gas density } n &\approx 2 \times 10^4 \text{ cm}^{-3}, \quad \text{wind velocity } v \approx 700 \text{ km s}^{-1}, \\ \text{mass - loss rate } \dot{M} &\approx 4 \times 10^{-3} \text{ M}_{\odot} \text{ yr}^{-1} \end{aligned} \tag{1}$$

(Geballe et al. 1987, 1991; Allen et al. 1990). If the wind consist mostly of helium, this can diminish n by a factor of two.

This outflow has been shown to arise from the sum of individual winds produced by hot stars of various locations and spectral types. They contain ~ 20 luminous blue supergiants (Krabbe et al. 1991), including several components of the central IRS 16 complex. Their P Cygni profiles directly indicate the existence of strong supersonic outflows (Najarro et al. 1994, Krabbe et al. 1995). The temperatures and spectral types of the above mentioned stars range from somewhat cooler, $\sim 20,000$ K blue supergiants with strong $\lambda 2.058 \mu\text{m}$ He I line emission to $\gtrsim 30,000$ K WN- and WC-type Wolf-Rayet (WR) stars. Mass loss rates of the individual stars range from a few 10^{-5} up to a few $10^{-4} \text{ M}_{\odot} \text{ yr}^{-1}$ at velocities of 300 to 900 km s^{-1} (Najarro et al. 1993, Krabbe et al. 1995, Blum et al. 1995, Tamblyn et al. 1996).

The presence of such powerful winds at the Galactic center has several profound implications for the central activity as well as for many morphological peculiarities found in the region:

- interception of part of the winds by a putative central black hole (Sgr A *) and the resulting wind accretion can serve as a source of the black hole feeding and accretion luminosity (Ozernoy 1989, 1993; Melia 1992; Narayan et al. 1995);
- interaction of the wind with the ionized and molecular gas at the Galactic center can explain much of its morphology (Yusef-Zadeh & Wardle 1993);
- both momentum and energy of the wind create a ‘wind-dominated’ regime within the central few tenths of a parsec, which influences substantially the dynamics of the region (Lutz et al. 1993).

These and other wind features are reviewed, in more detail and both from observational and theoretical sides, by Genzel, Hollenbach, & Townes (1994). Recently we have shown that the kinetic pressure of the wind is able to stop the inflow of matter into the central parsec from the circumnuclear ring and to reduce drastically the accretion rate onto the central black hole (Ozernoy & Genzel 1996, 1997). In the present paper, we would like to point out one more important corollary of the extended wind at the Galactic center – *collisions between the individual stellar winds within the core of the central stellar cluster*. The existence of a very dense core has been recently demonstrated by a high resolution imaging of the nuclear star

cluster (Eckart et al. 1993). Besides, both acceleration of particles by shock waves from the colliding winds in the cluster and hard X-ray and γ -ray emission of these particles are considered below.

2. COLLIDING WINDS IN THE STELLAR CORE

2.1. The physics of the collision and the production of X-rays

As is well known, among massive stars of WR and OB types, which have been found to be strong X-ray emitters, the most powerful are those belonging to binary systems. This phenomenon, commonly interpreted as due to the collisions of the winds from the components entering the binary, was at first predicted theoretically (Prilutskii & Usov 1976). These authors have shown that collision of the wind away from a WR star either with the wind from a companion OB star or with the surface of the latter induces a shock that heats the gas up to the temperature $T \sim 10^7$ K, resulting in X-ray emission with luminosity $L_X \sim 10^{33} - 10^{35}$ erg s $^{-1}$. By the present time, the theory of this process has been elaborated in much detail and it successfully serves to both interpret and guide the observations (for reviews, see e.g. Pollock 1987, 1995; Usov 1992, 1995; Stevens 1995).

It should be noted that the solitary massive OB stars also have inherent X-ray emission whose origin is related to shocks that appear as a result of instabilities in the radiation-driven winds from massive stars (Owocki & Rybicki 1985). However, OB stars belonging to binary systems are usually several times brighter at X-rays, and this has been explained by colliding winds (Chlebowski 1989). Moreover, the binary's L_X can be much higher if the companion of the OB star is a WR star rather than another OB star. This is due to the fact that the mass loss rate and the terminal velocity of the matter outflow far from the star are higher for WR stars

$$\dot{M}_{\text{WR}} \sim (0.8 - 8) \times 10^{-5} \text{ M}_{\odot} \text{ yr}^{-1}, \quad V_{\text{WR}}^{\infty} \sim (1 - 5) \times 10^3 \text{ km s}^{-1} \quad (2a)$$

than for OB stars of same masses

$$\dot{M}_{\text{OB}} \sim 10^{-6} \text{ M}_{\odot} \text{ yr}^{-1}, \quad V_{\text{OB}}^{\infty} \sim (1 - 3) \times 10^3 \text{ km s}^{-1} \quad (2b)$$

(e.g. Abbott & Conti 1987, Leitherer 1988).

In the case when both the WR and OB winds collide with their (supersonic) terminal velocities and X-ray absorption might be neglected, the resulting X-ray emission as a function of the binary's parameters is given by (Usov 1992):

$$L_X \simeq 10^{35} \left(\dot{M}_{-5}^{\text{WR}} \right)^{1/2} \left(\dot{M}_{-6}^{\text{OB}} \right)^{3/2} (V_3^{\text{WR}})^{1/2} (V_3^{\text{OB}})^{-3/2} D_{13}^{-1} \text{ erg s}^{-1}, \quad (3)$$

where $\dot{M}^{\text{WR}}$ and $\dot{M}^{\text{OB}}$ are normalized to $10^{-5} \text{ M}_{\odot} \text{ yr}^{-1}$ and $10^{-6} \text{ M}_{\odot} \text{ yr}^{-1}$, correspondingly; V^{WR} and V^{OB} are normalized to 10^3 km s^{-1} , and the distance D between WR and OB stars is normalized to 10^{13} cm . The radiation-driven coronal winds from massive stars attain their terminal velocity V^{∞} at the radius r_{ter} which is a few stellar photosphere radii (e.g. Friend & MacGregor 1984).

The analysis of the conditions under which the stellar winds collide in a WR + OB binary (Usov 1992) shows that neither a very close nor a very wide binary is able to produce strong X-ray emission. This is because in a very close binary the X-ray emission is strongly absorbed in the WR and OB winds (Luo, McCray & Mac Low 1990; Usov 1990; Stevens, Blondin & Pollock 1992), whereas in a very wide binary the X-ray luminosity decreases with an increase of the distance D between the components of the binary: $L_X \propto D^{-1}$, or even faster ($L_X \propto D^{-4/3}$) when no equalization of the ion and electron temperatures occurs within at least one shock layer around the contact surface (Usov 1992). It is expected that the X-ray luminosity of a WR + OB binary reaches its maximum, $L_X^{\max} \simeq 10^{35} \text{ erg s}^{-1}$, when $D \simeq \tilde{D} \equiv (\dot{M}_{\text{WR}} V_{\text{WR}}^\infty / \dot{M}_{\text{OB}} V_{\text{OB}}^\infty)^{1/2} r_{\text{ter}}$. For a typical WR + OB binary one gets $\tilde{D} \simeq 10^{13} \text{ cm}$ within a factor of 2-3 or so (Usov 1992).

So far massive binaries have been observed neither in IRS 16 nor elsewhere within the central parsec of the Galactic center, although lunar occultation observation do not exclude that some of the IRS 16 components might be multiple stars (Simons et al. 1990). Meanwhile all known massive stars within the central parsec are separated by such large distances ($\sim 10^{16} - 10^{17} \text{ cm}$) that the effects of their *individual* wind-wind collisions are rather weak (in contrast to the *collective* effects of wind collisions on the surrounding ISM – see Sec. 4). However, those effects could be much stronger, compared to IRS 16, in the central dense stellar cluster – the nuclear core. An inferred very high total stellar density in the core $\sim 10^7 - 10^8 \text{ stars/pc}^3$ (Eckart et al. 1993) implies comparatively frequent stellar encounters, with a part of them being as close as $\sim 10^{13} \text{ cm}$ to provide an effective collision of stellar winds and, as a result, the production of strong X-ray emission, acceleration of particles, and generation of related hard radiation (see below). An interesting circumstance is that we do not need to know an (unknown) number of WR + OB binaries: these close ‘pairs’ are formed temporarily during encounters in the cluster core to reveal this event by producing the above phenomena and then the stars fly apart not to meet each other anymore!

2.2. The rate of near-by stellar encounters in the stellar core

To calculate the rate of wind collisions within the stellar core, we introduce the number of wind producing stars, N_w . We assume that the radial distribution of these, most massive and luminous, stars follows an isothermal-like stellar distribution with a density run found by Eckart et al. (1993):

$$n(r) = n_c (r_c/r)^2, \quad (4)$$

where $r_c = 0.15 \pm 0.05 \text{ pc}$ is the core radius, $\rho_c = 9\sigma_v^2/4\pi G r_c^2 = 10^{7.9 \pm 0.5} \text{ M}_\odot \text{ pc}^{-3}$ is the inferred mass density within the core, and $\sigma_v = 100 \pm 25 \text{ km s}^{-1}$ is the line-of-sight central velocity dispersion. In fact, available data indicate (Eckart et al. 1993, Genzel et al. 1996) that the early type stars may have even a slightly smaller core radius (0.1 pc) whereas for the late type stars it is larger ($r_c \sim 0.3 - 0.5 \text{ pc}$).

While calculating the rate with which the winds in the core collide, we constrain ourselves only with those encounters of the wind producing stars whose collision

periastron is $D \lesssim \tilde{D} \simeq 10^{13}$ cm. The rate of stellar encounters with an impact parameter D occurring in an isothermal spherical cluster is given by

$$\mathcal{R} \simeq 20\pi^{3/2} n_w^2 r_c^3 \left[4D^2 \sigma_v + \frac{G(m_{w1} + m_{w2})D}{\sigma_v} \right], \quad (5)$$

where the second term in the brackets accounts for the gravitational focusing of the encountering stars whose masses are m_{w1} and m_{w2} .

In order to evaluate a characteristic value of $\mathcal{R}$, let us take $m_{w1} = m_{w2} = 10 M_\odot$ and put $n_w \equiv 10^7 n_7$ stars/pc³ and $r_c \equiv 0.1 r_{-1}$ pc. One gets

$$\mathcal{R} \simeq 3.8 \cdot 10^{-10} n_7^2 r_{-1}^3 \left[4D_{13}^2 \sigma_2 + 2.7 \frac{D_{13}}{\sigma_2} \right] \text{ s}^{-1} \sim 0.1 n_7^2 r_{-1}^3 \text{ yr}^{-1}, \quad (6)$$

where $\sigma_2 \equiv \sigma_v/100 \text{ km s}^{-1}$. Given the number of stars producing winds N_w , their density within the core is $n_w = N_w / (\frac{4}{3}\pi r_c^3) \simeq 2.4 \cdot 10^5 N_3 r_{-1}^{-3} \text{ pc}^{-3}$, where $N_3 \equiv N_w/10^3$. Eq. (6) yields then

$$\mathcal{R} \sim 0.6 \cdot 10^{-4} N_3^2 r_{-1}^{-3} \text{ yr}^{-1}. \quad (7)$$

2.3. Parameters of X-ray flares due to colliding winds

To evaluate an actual collision rate, one needs to know the number of wind producing stars within the core. Evidently, early-type stars would be of prime interest here. Let us discuss available constraints to the number of those stars.

The entire bolometric luminosity within $r \leq 0.6$ pc at the Galactic center is $L_{\text{bol}} \simeq 2 \times 10^7 L_\odot$, with about 10% of it in the form of Ly α photons (Zylka et al. 1995, Krabbe et al. 1995). The luminosity as high as this could be provided by just $\sim 10^2$ luminous stars like the Allen-Forrest (AF) object (Allen et al. 1990, Forrest et al. 1987) with $L_{\text{AF}} \simeq (2 - 3) \times 10^5 L_\odot$ and mass-loss rate $\dot{M} \simeq (6 \pm 1) \times 10^{-5} M_\odot \text{ yr}^{-1}$ (Najarro et al. 1994). However the stars as luminous as the AF star are rather rare.

Let us suppose, for a moment, that the ‘wind-dominating’ region at the Galactic center is created by much more numerous, although less luminous but still massive, stars. Using as a scaling an empirical relationship $\dot{M} \propto L^{1.25}$ (van Buren 1985) and normalizing it with the help of the AF star, one can see that both the total mass-loss rate $\dot{M} \simeq 4 \times 10^{-3} M_\odot \text{ yr}^{-1}$ [Eq. (1)] and the total luminosity of the wind producing region could be reproduced by N_w stars of luminosity L_w and mass-loss rate $\dot{M}_w$ each. The values of N_w , L_w , and $\dot{M}_w$ are presented in Table 1:

% PUT TABLE 1 HERE %

As Eq. (7) shows, even if the number of massive, wind producing stars in the core were as large as $N_w = 10^4$, the rate of near, $D \lesssim 10^{13}$ cm, stellar encounters would only be $\mathcal{R} \sim 0.6 \times 10^{-2} \text{ yr}^{-1}$. Besides, those winds would have $\dot{M} \simeq 2 \times 10^{-7} M_\odot \text{ yr}^{-1}$, at which according to Eq. (3) their collisional $L_X \ll 10^{35} \text{ erg/s}$. Only a combination of $N_w \lesssim 10^3$ and $r_c \ll 0.1$ pc would make the dissipation of

kinetic energy of the colliding winds observable as X-ray flares with both $L_X \sim 10^{35}$ erg s $^{-1}$ and high enough rate of these flares. The duration of the flares is

$$\Delta t \simeq \frac{D}{\sqrt{3}\sigma_v} \sim 1 \frac{D_{13}}{\sigma_2} \text{ week.} \quad (8)$$

Taking into account that $L_X \propto D^{-1}$ (see Eq. (3)), Eqs (6) and (8) yield $\mathcal{R} \propto (\Delta L_X)^{-2}$ and $\Delta t \propto (\Delta L_X)^{-1}$, where ΔL_X is the amplitude of modulations of the X-ray luminosity of the compact cluster at the Galactic center because of the wind collisions. For $N_w = 10^3$, we have numerically $\mathcal{R} \sim 10^{-4}(\Delta L_X/10^{35} \text{ erg s}^{-1})^{-2} r_{-1}^{-3} \text{ yr}^{-1}$ and $\Delta t \sim 1(\Delta L_X/10^{35} \text{ erg s}^{-1})^{-1} \sigma_2^{-1} \text{ week}$. From both the condition $\mathcal{R} \simeq (\Delta t)^{-1}$ and the fact that the soft X-ray luminosity of the Galactic Center is $\sim 1.5 \times 10^{35} \text{ erg s}^{-1}$, the X-ray flux from the Galactic Center has to be variable on the time scale of one year with the amplitude of $\sim 1 \%$.

Available high resolution near-infrared imaging data obtained with the MPE camera SHARP (Eckart et al. 1993, 1995) enable us, from counting the number of objects on the SHARP map, to derive an upper limit for the number of stars earlier than B0.5/B1 [$K \sim 15$, $L \sim (5 - 10) \times 10^3 L_\odot$]. Taking into account the uncertainty due to confusion, noise, and completeness at $K \sim 15$, there are no more than ~ 100 to 200 such stars within $r \lesssim 4''$ of the central stellar core. Having compared this upper limit with Eq. (7) one concludes that detections of individual X-ray flares from collisions of stellar winds in the dense core are untenable at the Galactic center.

The rate of wind collisions could be substantially increased if one accounts for the influence of a putative massive black hole thought to present at the Galactic center as a compact radio source Sgr A * . The rate of stellar collisions in a density spike around Sgr A * is enhanced by a factor of 30-40 (Ozernoy 1994). Even if the density spike has already been destroyed by those frequent stellar collisions the gravitational field of the black hole enhances stellar density of unbound stars in the black hole vicinity and hence increases appreciably the rate of wind collisions. Therefore, X-ray flares from the colliding winds at the Galactic center, if detected, would provide an indirect evidence in favor a massive black hole sitting there.

Unless the site for colliding stellar winds is more compact and/or a massive black hole enhances the collision rate in its vicinity, a larger number of wind producing stars would result in an observable effect. Yet the disentangling of the thermal, wind-produced X-ray flares from the persistent X-ray emission turns out to be rather difficult partly because that emission is due to the (shocked) global wind itself. Meanwhile the above consideration, exemplified by the Galactic center case, reveals conditions under which the thermal X-ray flares produced by colliding winds could be observable in some compact star clusters elsewhere. In the Galactic center, the overall soft X-ray emission due to the shocked winds in the central cavity ($R \approx 1.5$ pc) is given by $L \approx 10^{35} (p_{th}/10^8 \text{ cm}^{-3} \text{ K})^2 f_V \text{ erg s}^{-1}$ (Genzel et al. 1994), where $p_{th} \approx 10^8 \text{ cm}^{-3} \text{ K}$ is the characteristic thermal pressure in the hot ($T \simeq 10^7 \text{ K}$) shocked wind and f_V is the volume filling factor of the shocked gas. Therefore, the persistent emission would be temporally modulated by individual X-ray flares if the (otherwise similar) star cluster had a smaller p_{th} implying more rarefied gas.

3. NON-THERMAL EMISSION OF PARTICLES ACCELERATED IN SHOCKS FROM COLLIDING WINDS

As is known, colliding winds which flow away from massive stars are able to accelerate charged particles via shock waves. Indeed, several WR + OB binaries demonstrate non-thermal radio emission that has been shown to be associated with the accelerated particles (Eichler & Usov 1993 and references therein). Spectral properties of that radio emission are consistent with what would be expected from the electrons which are accelerated by the shocks produced by colliding winds (Eichler and Usov, 1993; Dougherty et al. 1996).

3.1. Conditions for particle acceleration and the outflowing gas structure

To accelerate particles by shocks, the strength of the magnetic field at the region of stellar wind collision, at the distance of a few 10^{16} cm from a WR star, has to be within the range $10^{-12} \ll B \ll 3 \times 10^{-3}$ Gauss [see eqs. (8)-(12) in Eichler and Usov 1993]. The strength of the magnetic field in the plasma flowing away from a WR star is given, far enough from the star ($r > RV_{WR}^\infty/V_{\text{rot}}$), by

$$B \simeq B_s \frac{V_{\text{rot}}}{V_{WR}^\infty} \frac{R^2}{r_A r} \quad (9)$$

(e.g., Usov and Melrose 1992), where B_s is the surface magnetic field, R is the stellar radius, V_{rot} is the surface rotational velocity, r_A is the Alfven radius, and V_{WR}^∞ is the terminal velocity of the WR wind. For the fiducial values of $B_s \sim 10^2$ G, $R \sim 10^{12}$ cm, and $V_{\text{rot}} \sim 0.1V_{WR}^\infty$ at $r \sim \text{a few} \times 10^{16}$ cm, one gets $B \sim \text{a few} \times 10^{-4}$ G, which is well within the range of B given above. This and the other conditions which are necessary for particle acceleration by shocks (e.g., Eichler & Usov, 1993) are met in the colliding winds at the stellar core at the Galactic Center. Therefore, charged particles (electrons, protons, and nuclei) can be accelerated in the core to high enough energies. The maximum Lorentz factor of accelerated electrons is about 10^5 [see eqs. (13)-(14) in Eichler and Usov (1993)].

The colliding winds outflowing from the massive stars in the central stellar cluster produce an ensemble of shock waves. Indeed, although both the structure and kinematics of the interstellar matter inside the stellar cluster are rather complex, the structure of that gas can be considered as a network of N_w bubbles, or ‘cells’ formed around the wind-producing stars. The mean size of the cell is about $l_w \simeq lN_w^{-1/3}$. Inside each cell, the wind gas is comparatively cold ($\sim 10^4$ K) and moves away from the star until it collides with the ambient gas. The latter, being a mixture of the stellar winds outflowing from the massive stars, is rather hot ($\sim 10^6 - 10^7$ K, see Sect.4) due to collisions of many individual stellar winds. Whenever stellar wind from a massive star collides with the ambient gas, a shock wave is formed in the wind and the outflowing gas is heated. The shocked gas joins the hot ambient gas and moves away from the central cluster through the labyrinth formed by the network of cells. The walls of this labyrinth are the shock fronts. Since the wind velocities exceed, by a factor of ten, the velocity dispersion of massive stars, the gas flow in the

central cluster can be considered as quasi-stationary. The structure of the interstellar matter inside the stellar cluster varies on a time scale of $\sim l_w/\sigma_v \sim 30$ yr.

3.2. Particle acceleration by multiple shocks and radiation of the accelerated particles

Multiple shocks produced by the colliding winds accelerate particles to high energies until the particles escape from the cluster. Diffusive acceleration of particles by multiple shocks produces a distribution in momentum, which is flatter than that produced by a single shock (e.g., Melrose and Pope 1993); specifically, the distribution approaches $f(p) \propto p^{-b}$, with $b = 3$ for a large number of strong shocks, while $b = 4$ for a single strong shock. A synchrotron spectrum $I_\nu \propto \nu^{-\alpha}$ with $\alpha = 0$ corresponds to the distribution function of relativistic electrons with $b = 3$. The following simplifications have been made to get this asymptotical distribution of accelerated particles: (i) all shocks are identical, (ii) decompression occurs between each shock, and (iii) at each shock an injection of new particles takes place. In a real situation, the spectral index may differ from $b = 3$. Yet, it is natural to expect that the synchrotron spectrum of high energy electrons accelerated by an ensemble of shocks is more or less flat, i.e. $\alpha \simeq 0$.

Synchrotron radiation of high energy electrons produced in the cluster has to be suppressed, owing to the Razin-Tsytovich effect (Tsytovich 1951; Razin 1960), at low frequencies $\nu < \nu_{\text{RT}}$ such that

$$\nu_{\text{RT}} \simeq 20(n_e/B) \text{ Hz}, \quad (10)$$

where the electron number density, n_e , is given in cm^{-3} , and B in Gauss. For $n_e \simeq 2 \times 10^4 \text{ cm}^{-3}$ and $B \simeq 2 \times 10^{-4} \text{ Gauss}$ this yields $\nu_{\text{RT}} \simeq 2 \text{ GHz}$.

The expected synchrotron luminosity at $\nu > \nu_{\text{RT}}$ is given by

$$L_s \simeq \xi \frac{\tau_{\text{esc}}}{\tau_s} L_{\text{wind}}, \quad (11)$$

where ξ is the efficiency of electron acceleration by shocks, $L_{\text{wind}} = \frac{1}{2} \dot{M} v^2 \simeq 3 \times 10^{38} \text{ ergs s}^{-1}$, $\tau_{\text{esc}} \simeq l/v \simeq 4 \times 10^9 \text{ s}$ is the characteristic time of escape of high energy electrons from the cluster, and

$$\tau_s \simeq \frac{5 \times 10^8}{B^2 \gamma_e} \text{ s} \quad (12)$$

is the characteristic time of synchrotron losses by relativistic electrons of the Lorentz factor γ_e . Since the Lorentz factor of high energy electrons, whose synchrotron emission in the field B has a maximum at the frequency ν , is given by

$$\gamma_e \simeq 2 \times 10^3 \left(\frac{B}{10^{-4} \text{ Gauss}} \right)^{-1/2} \left(\frac{\nu}{1 \text{ GHz}} \right)^{1/2}, \quad (13)$$

one gets from eqs. (11)-(13), for $B \simeq 2 \times 10^{-4} \text{ Gauss}$, $\xi \simeq 0.1$, and $\nu \simeq 5 \text{ GHz}$ that $L_s \simeq 3 \times 10^{34} \text{ erg s}^{-1}$, which is comparable to the total radio luminosity of

the compact radio source Sgr A*. To be more accurate, the value of L_s should be considered as an upper limit to the synchrotron emission of high energy electrons, which are accelerated by an ensemble of shocks in the stellar cluster.

Electrons that are accelerated in the core can serve as a source of non-thermal X-ray and γ -ray emissions due to the Compton scattering of the respective IR and UV photons off the electrons. The spectrum of those emissions is expected to be flat, like the synchrotron spectrum discussed above. The non-thermal luminosity due to the scattered photons is roughly given by

$$L_c \simeq \left(\frac{\epsilon_{\text{ph}}}{B^2/8\pi} \right) L_s, \quad (14)$$

where

$$\epsilon_{\text{ph}} \simeq \frac{L_{\text{IR+UV}}}{4\pi r_c^2 c} \quad (15)$$

is the photon energy density in IR and UV ranges inside the cluster, and $L_{\text{IR+UV}}$ is the total IR + UV luminosity of the cluster. Certainly, the value of L_c cannot exceed $\xi L_{\text{wind}} \simeq \text{a few } \times 10^{37} \text{ erg s}^{-1}$. For the same parameters of the cluster and $L_{\text{IR+UV}} \simeq 10^{40} \text{ erg s}^{-1}$, eqs. (14) and (15) yield $L_c \simeq 5 \times 10^{36} \text{ erg s}^{-1}$. Compton scattering of the photons having energies $\sim 0.1 - 1 \text{ eV}$ off the electrons having $\gamma_e \sim 10^3$ produces the photons of the mean energy about $10^2 - 10^3 \text{ MeV}$, which is in the *EGRET* range of the *Compton Gamma Ray Observatory*. Our estimated value of L_c is comparable to, but somewhat below than, the observed value of $L_\gamma \sim 10^{37} \text{ erg s}^{-1}$ (Mattox et al. 1993, Mayer-Hasselwander et al. 1993).

Shocks from the colliding winds in the stellar cluster at the Galactic center can accelerate, besides electrons, protons and nuclei as well. The decay of pions that emerge from baryon – baryon collisions results in the production of gamma-rays. Before escaping from the stellar cluster, relativistic nuclei pass through the gas of column density $\Sigma_r \sim n(l/v)c$. Putting $n \simeq 2 \times 10^4 \text{ cm}^{-3}$, $l \simeq 0.1 \text{ pc}$, and $v \simeq 700 \text{ km s}^{-1}$, one gets $\Sigma_r \sim 10^{24} \text{ cm}^{-2}$. The cross section of π^0 -meson production is about $3 \times 10^{-26} \text{ cm}^2$ for colliding protons near the threshold of this process, i.e. at $E_p \sim 0.3 - 1 \text{ GeV}$, and is larger for nuclei. Therefore, for $\Sigma_r \sim 10^{24} \text{ cm}^{-2}$ the efficiency of pion production is somewhat between a few % and 10%. Such an efficiency seems plausible for conversion of the energy of accelerated protons and nuclei into the energy of γ -rays.

Before going out from the stellar core, the outflowing gas has to pass through one or more shocks. Therefore, one can expect that the luminosity of the core in cosmic rays might be as high as

$$L_{CR} \sim 0.1 L_{\text{wind}} \sim 3 \times 10^{37} \text{ erg s}^{-1}. \quad (16)$$

The luminosity in γ -rays owing to pion decay can reach $L_\pi \sim 0.1 L_{CR} \sim \text{a few } \times 10^{36} \text{ ergs s}^{-1}$. For γ -rays from pion decays, there should be a low-energy spectral cutoff at the energy $E_{\gamma,c} \sim 70 \text{ MeV}$. At energies $E \gg E_{\gamma,c}$, the spectral index of γ -rays is expected to be similar to the spectral index of accelerated nuclei. The total

luminosity of the core in γ -rays might be as high as $\sim 10^{37}$ ergs s $^{-1}$ within a factor of 2 or so. The spectral observations of γ -rays from the core could distinguish between the inverse Compton γ -ray emission and γ -rays generated by pion decays. This would allow to test the above picture and to estimate the model parameters.

4. DISCUSSION

Colliding winds in the core of the Galactic center's stellar cluster have two basic differences from those in massive WR + OB binaries. First, one cannot expect to keep such binaries in the core till the present time [because in all 'relic' massive binaries which might have formed during the latest starburst in the Galactic nucleus 5-8 Myr ago, at least one component has already evolved into a compact remnant (Lipunov et al. 1996), whereas formation and hardening of new hard binaries via triple or double encounters of massive stars would require a time that exceeds much the life-time of those stars on the main sequence]. Nevertheless, a very high stellar density in the core makes it feasible for the encounters between the wind producing stars to be so close as to provide strong collisions between their winds. Second, the winds from individual stars in the cluster core propagate through a rarefied, hot ambient gas that is produced by a superposition of many individual stellar winds.

Although this environment looks different from that in massive binaries, it does not influence the conditions under which, at a close encounter of two stars in the core, their winds collide so as to produce a strong X-ray emission: Indeed, the termination shock radius, R_s , relative to an individual star with mass-loss rate $\dot{M}_w$ and velocity of the wind V_w is given by the condition that the pressure downstream from the shock equals the sum of thermal and ram pressures of the ambient gas, i.e.,

$$\frac{\dot{M}_w V_w}{4\pi R_s^2} = p_{th} + \rho v^2. \quad (17)$$

The central value of the pressure in the stellar core is of thermal origin but in the bulk of the core and elsewhere in the central parsec the ram pressure dominates the pressure of the outflowing gas. At a characteristic radius r from the center of the core, the distance of the termination shock from the star is found to be $R_s \approx \left(\dot{M}_w V_w / \dot{M} v \right)^{1/2} r$. At $\dot{M}_w = 10^{-6} M_\odot \text{ yr}^{-1}$, $V_w^\infty = 10^3 \text{ km s}^{-1}$, and $\dot{M}$ and v given by Eq. (1), this yields $R_s \approx 0.01 \text{ pc}$ at $r \simeq 0.3 \text{ pc}$. Thus R_s is much larger than the characteristic distance of 10^{13} cm , at which one expects the X-ray luminosity that is about the total soft X-ray luminosity of the Galactic center. In other words, the ambient ram pressure is not too large to have any influence on the colliding winds when the impact parameter is $D \lesssim 10^{15} \text{ cm}$.

At the same time, the ambient ram pressure is large enough to provide a high dynamic importance in the entire region occupied by the shocked winds. As a particular example, let us consider the origin of the wind thought to be responsible for the dynamics of the Galactic center 'mini-cavity' (for a review of the 'mini-cavity' 's morphology and dynamics, see Genzel et al. 1994). Several hypotheses have been explored: (i) winds from IRS 16 complex are deflected and collimated by

the gravitational field of Sgr A* (Wardle & Yusef-Zadeh 1992), (ii) winds due to one or several stars very close to the center of the mini-cavity (Lutz et al. 1993), and (iii) a wind that comes directly from Sgr A* in the form of a jet (Lutz et al. 1993). A recent finding that Sgr A* lies very close to or even within the core of the central stellar cluster (Eckart et al. 1993) and an enhanced density of stellar winds in the cluster core both enable us to explore one more possibility: the wind that provides the energetics of the ‘mini-cavity’ originates in the cluster core. A presence within the core of even $N_w = 10$ wind producing stars with a terminal outflow velocity $V_w^\infty = 10^3 \text{ km s}^{-1}$ and an outflow rate $\dot{M}_w = 10^{-5} \text{ M}_\odot \text{ yr}^{-1}$ each (or $N_w = 100$ with $\dot{M}_w = 10^{-6} \text{ M}_\odot \text{ yr}^{-1}$) could provide a total wind mechanical luminosity of about $3 \times 10^{37} \text{ erg s}^{-1}$, which is consistent with the estimate that follows from the dynamics of a spherically-symmetric wind bubble (Lutz et al. 1993). However, an off-center location of the ‘mini-cavity’ does not easily fit in with a model of a symmetric wind source centered on the dynamic center. Furthermore, if N_w is too large, that would change the present morphology of the ‘mini-cavity’ and induce a similar cavity elsewhere. Therefore, the ‘mini-cavity’ phenomenon could be used for further constraining the number of wind-producing stars in the core.

Collisions of hot winds at the Galactic center result in (i) formation of an ensemble of shock waves, (ii) particle acceleration by the shocks, and (iii) generation of hard X-ray and γ -ray emission just as in massive binaries. Although neither the origin nor the total number of massive, wind producing stars in the cluster core is known as yet, the production of a strong emission by the colliding winds (Sec. 3) could, in principle, be used to constrain the value of N_w . However, as is shown in Sec. 2, the rate of wind collisions able to produce each $L_X \simeq 10^{35} \text{ erg s}^{-1}$ is only $\mathcal{R} \sim 10^{-4} \text{ yr}^{-1}$ if $N_w \sim 10^3$. Accounting for the influence of a putative massive black hole thought to present at the Galactic center as a compact radio source Sgr A* would increase this rate by a factor of 30-40 if the black hole mass is as large as $M_h \sim 10^6 \text{ M}_\odot$. We also note in passing that the clumps produced by the colliding winds being accreted onto the black hole would have induced a strong flaring activity of Sgr A*. The recent finding of a clustering of luminous stars in the immediate vicinity of Sgr A* (Eckart et al. 1995) might indicate an enhanced rate of wind collisions close to this enigmatic source.

Let us discuss the expected hard radiation resulting from acceleration of particles by the shock-wave ensemble in the cluster of massive stars at the Galactic center. As is shown in Sect. 3, acceleration of particles at shocks produced by the colliding winds at the Galactic center’s stellar core, might result in continuum γ -ray emission as high as $L_\gamma \sim 10^{37} \text{ erg/s}$. Besides, some γ -ray lines could be expected as well. To evaluate the intensity of that emission, it is helpful to address to the γ -ray line emission from the Orion complex recently claimed to be detected in the 3-7 MeV range (Bloemen et al. 1994). Those γ -ray lines have been interpreted as the de-excitation lines $^{12}\text{C}^*(4.43 \text{ MeV})$ and $^{16}\text{O}^*(6.13 \text{ MeV})$ due to either cosmic-ray carbon and oxygen nuclei interacting with the ambient medium (Bloemen et al. 1994, Bykov & Bloemen 1994) or a combination of those cosmic-ray nuclei plus cosmic-ray protons and α -particles interacting with the ambient C and O nuclei (Ramaty et al. 1995). Although a recent re-evaluation of the data indicates that their

original interpretation meets serious problems (Bloemen et al. 1997), it is instructive to estimate the expected γ -ray emission associated with the colliding winds at the Galactic center and then to compare it with the Orion data. The column density of gas in the stellar core at the Galactic center is estimated from Eq. (1) to be $\Sigma \simeq 0.6 \times 10^{22} \text{ cm}^{-2}$, which is larger, by a factor of 10^2 , than that in the Orion complex as given by Cowsik & Friedlander (1995). Our upper limit to the number of early-type stars in the vicinity of the Galactic center's stellar core ($N_w < 100$ -200, see Sect.2.3) is larger than (although comparable to) the number of hot massive stars in the Orion ($N = 56$, according to Genzel & Stutzki 1989). Accordingly, the total kinetic luminosity of the winds from massive stars in the core is more than that in the Orion complex. (The expected luminosity in γ -rays is proportional to the product of ΣL_w .) Since the distance to the Galactic center exceeds that to the Orion complex (about 450 pc) by a factor of 20, we expect that the flux of γ -rays from the Galactic center in the 3-7 MeV range is comparable to that from the Orion complex. This conclusion is consistent with the *SMM* measurements (Harris et al. 1995) that the total Galactic center (from its central radian) γ -ray flux is no greater than 4.7 times the published *COMPTEL* flux from Orion at 4.4 MeV. Therefore, γ -rays from the Galactic center's stellar core might be a target to search for in further missions such as *INTEGRAL*. Detecting or constraining that γ -ray emission, including the line emission of heavy nuclei such as carbon and oxygen, at 4.4 and 6.1 MeV, would be important to further constrain the parameters of massive stars in the cluster at the Galactic center.

Finally, it is worth mentioning that the winds colliding at comparatively large impact parameters are producing rather strong radio emission. It has recently been observed that in the vicinity of Sgr A* there are blobs of ionized plasma detected by their radio emission and claimed to be either ejected from this source (Yusef-Zadeh et al. 1990) or resulted from a not yet specified interaction of the global wind with the gravitational potential of Sgr A* (Wardle & Yusef-Zadeh 1992). One more possibility that needs to be explored in more detail is that those blobs could be the products of the colliding winds in the vicinity of Sgr A*. The inferred parameters of the blobs, $n \simeq 10^4 \text{ cm}^{-3}$, $T \simeq 10^4 \text{ K}$ (Yusef-Zadeh et al. 1990), imply that these blobs are already in the thermal equilibrium with the shocked wind, $p_{th} \sim 10^8 \text{ cm}^{-3} \text{ K}$ (Genzel et al. 1994).

Among those observations which could make further progress in our understanding of the origin of the wind in Sgr A*/IRS 16 complex, we would like to mention subarcsecond radio continuum maps of mass-losing stars in the vicinity of Sgr A*. Such observations would help to resolve some problems related to the origin of the wind, including the type of wind producing stars. The standard model for a spherically symmetric wind of partially ionized gas (molecular weight μ , average ionic charge $\bar{Z}$) with $T \simeq 10^4 \text{ K}$ and $\dot{M} \simeq 6 \times 10^{-5} M_\odot \text{ yr}^{-1}$ expanding with a constant velocity v would give the flux density of free-free radio emission

$$F_\nu \approx 5 \text{ mJy} \left(\frac{\mu}{1.2} \right)^{-4/3} \bar{Z}^{2/3} \left(\frac{\nu}{100 \text{ GHz}} \right)^{0.6} \left(\frac{\dot{M}}{6 \times 10^{-5} M_\odot \text{ yr}^{-1}} \right)^{4/3}$$

$$\times \left(\frac{v}{700 \text{ km s}^{-1}} \right)^{-4/3} \left(\frac{d}{8.5 \text{ kpc}} \right)^{-2} \quad (18)$$

(Panagia and Felli 1975; Wright and Barlow 1975). Measuring F_ν at millimeter wavelengths with VLA interferometry would yield $\dot{M}$, which with the use of the empirical relationship $\dot{M} \propto L^{1.25}$ (van Buren 1985) would give L and therefore important information about the spectral types of stars responsible for individual winds.

Acknowledgements

We are grateful to an anonymous referee for helpful comments.

Table 1
Luminosity and Mass-loss Rate of an Average Wind Producing Star

Table 1

N_w	$L/L_\odot$	$\dot{M}_w/M_\odot\text{yr}^{-1}$
10^2	$3 \cdot 10^5$	$6 \cdot 10^{-5}$
10^3	$3 \cdot 10^4$	$3 \cdot 10^{-6}$
10^4	$3 \cdot 10^3$	$2 \cdot 10^{-7}$

REFERENCES

- Abbott, D.C. & Conti, P.C. 1987, ARA&A 25, 113
- Allen, D.A., Hyland, A.R., & Hillier, D.J. 1990, MNRAS, 244, 706
- Bloemen, H. 1996 (to be published)
- Bloemen, H. et al. 1994, Astron. Astrophys. 281, L 5
- Bloemen, H. et al. 1997, ApJ 475, L25
- Blum, R.D., Sellgren, K., & DePoy, D.L. 1995, ApJ 440, L 17
- Bykov, A. & Bloemen, H. 1994, Astron. Astrophys. 283, L1
- Chlebowski, T. 1989, ApJ 342, 1091
- Cowsik, R. & Friedlander, M.W. 1995, ApJ 444, L 29
- Dougherty, S.M., Williams, P.W., van der Hucht, K.A., Bode, M.F. & Davis, R.J. 1996, MNRAS 280, 963
- Eckart, A., Genzel, R., Hoffmann, R., Sams, B.J., & Tacconi-Garman, L.E. 1993, ApJ, 407, L 77
- Eckart, A., Genzel, R., Hoffmann, R., Sams, B.J., & Tacconi-Garman, L.E. 1995, ApJ, 445, L 23
- Eichler, D. & Usov, V.V. 1993, ApJ, 402, 27
- Forrest, W.J., Shure, M.A., Pipher, J.L., & Woodward, C.A. 1987, in “The Galactic Center”, ed. D.C. Backer. AIP Conf. Proc. 155, 153
- Friend, D.B. & MacGregor, K.B. 1984, ApJ 282, 591
- Geballe, T.R., Persson, S.E., Lacy, J.H., Neugebauer, G., & Beck, S.C. 1982, in “The Galactic Center”, ed. G.R. Riegler & R.D. Blandford (New York: Am. Inst. Phys.), p. 60
- Geballe, T.R., Krisciunas, K.L., Lee, T.J., Gatley, I., Wade, R., Duncan, W.D., Garden, R., & Becklin, E.E. 1984, ApJ, 284, 118
- Geballe, T.R., Wade, R., Krisciunas, K.L., Gatley, R., & Bird, M.C. 1987, ApJ, 320, 562
- Geballe, T.R., Krisciunas, K., Bailey, J.A., & Wade, R. 1991, ApJ 370, L 73
- Genzel, R., Hollenbach, D., & Townes, C.H. 1994, Rep. Progr. Phys. 57, 417
- Genzel, R. & Stutzki, J. 1989, ARA&A 27, 41
- Genzel, R., Thatte, N., Krabbe, A., Kroker, H., Tacconi-Garman, L.E. 1996, MPE Preprint 362=ApJ (submitted)
- Hall, D.N.B., Kleinmann, S.G., & Scoville, N.Z. 1982, ApJ 262, L 53
- Harris, M.J., Share, G.H., & Messina, D.C. 1995, ApJ 448, 157
- Krabbe, A., Genzel, R., Drapatz, S., & Rotaciuc, V. 1991, ApJ 382, L 19
- Krabbe, A., Genzel, R., Eckart, A., Naharro, F., Lutz, D., Cameron, M., Kroker, H., Tacconi-Garman, L.E., Thatte, N., Weitzel, L., Drapatz, S., Geballe, T., Sternberg, A., & Kudritzki, R. 1995, ApJ 447, L 95
- Leitherer, C. 1988, ApJ 326, 356
- Lipunov, V.M., Ozernoy, L.M., Popov, S.V., Postnov, K.A., & Prokhorov, M.E. 1996, ApJ 466, 234
- Luo, D., McCray, R., & Mac Low, M.-M. 1990, ApJ 362, 267
- Lutz, D., Krabbe, A., & Genzel, R. 1993, ApJ 418, 214
- Mattox, J.R. et al. 1993, BAAS 24, 1296
- Mayer-Hasselwander, H.A. et al. 1993, Proc. 23rd Intern. Cosmic Ray Conf. 1, 147

- Melia, F. 1992, *ApJ*, 387, L25
- Melrose, D.B. & Pope, M.H. *Proceedings ASA* 10, 222, 1993
- Najarro, F., Hillier, D.J., Kudritzki, R.P., Krabbe, A., Lutz, D., Genzel, R., Drapatz, S., & Geballe, T.R. 1994, *A&A* 285, 573
- Narayan, R., Yi, I., & Mahadevan, R. 1995, *Nat* 374, 623
- Owocki, S.P. & Rybicki, G.B. 1985, *ApJ* 299, 265
- Ozernoy, L.M. 1989, in “The Galactic Center”, *Proc. IAU Symp. No. 136*, ed. M. Morris (Dordrecht: Reidel), p. 555
- Ozernoy, L.M. 1993, in “Back to the Galaxy”, eds. S. Holt & F. Verter. *AIP Conf. Proc.*, 278, 69
- Ozernoy, L.M. 1994, in “Nuclei of Nearby Galaxies: Lessons from the Galactic Center”, eds. R. Genzel and A. Harris (Kluwer Acad. Publ.), p. 431
- Ozernoy, L.M. & Genzel, R. 1996a, in “The Unsolved Problems of the Milky Way”, *Proc. IAU Symp. No. 169*, ed. L. Blitz & P. Teuben (Kluwer Acad. Publ.), p. 181
- Ozernoy, L.M. & Genzel, R. 1996b, to be submitted
- Panagia, N. & Felli, M. 1975, *A&A*, 39, 1
- Pollock, A.M. 1987, *ApJ* 320, 283
- Pollock, A.M. 1995, in “Wolf-Rayet Stars: Binaries, Colliding Winds, Evolution”, *Proc. IAU Symp. No. 163*, eds. K.A. van der Hucht & P.M. Williams (Kluwer Acad. Publ.), p. 429
- Prilutskii, O.F. & Usov, V.V. 1976, *Sov. Astr.* 20, 2
- Ramaty, R., Kozlovsky, B., & Lingenfelter, R. 1995, *ApJ* 438, L 21
- Razin, V.A. 1960, *Radiofizika* 3, 584
- Stevens, I.R. 1995, in “Wolf-Rayet Stars: Binaries, Colliding Winds, Evolution”, *Proc. IAU Symp. No. 163*, eds. K.A. van der Hucht & P.M. Williams (Kluwer Acad. Publ.), p. 486
- Stevens, I.R., Blondin, J.M., & Pollock, A.M.T. 1992, *ApJ* 386, 265
- Tamblyn, P., Rieke, G.H., Hanson, M.M., Close, L.M., McCarthy, D.W., & Rieke, M.J. 1996, *ApJ* 456, 206
- Tsyтович, V.N. 1951, *Vest. Mosk. Univ. Phys.* 11, 27
- Usov, V.V. 1990, *Ap&SS* 167, 297
- Usov, V.V. 1992, *ApJ* 389, 635
- Usov, V.V. 1995, in “Wolf-Rayet Stars: Binaries, Colliding Winds, Evolution”, *Proc. IAU Symp. No. 163*, eds. K.A. van der Hucht & P.M. Williams (Kluwer Acad. Publ.), p. 495
- Usov, V.V. & Melrose, D.B. 1992, *ApJ*, 395, 575
- van Buren, D. 1985, *ApJ*, 294, 567
- Wardle, M. & Yusef-Zadeh, F. 1992, *Nat* 357, 308
- Wright, A.E., & Barlow, M.J. 1975, in “The Galactic Center”, *Proc. IAU Symp. No. 136*, ed. M. Morris (Dordrecht: Reidel), p. 443
- Yusef-Zadeh, F., Morris, M., & Ekers, R.D. 1990, *BAAS*, 22, 865
- Yusef-Zadeh, F. & Wardle, M. 1993, *ApJ*, 405, 584
- Zylka, R., Mezger, P.G., Ward-Thompson, D., Duschl, W.J. & Lesh, H. 1995, *AA*, 297, 83